

Variant: *NM_000173.7(GP1BA):c.673T>C (p.Cys225Arg)*

Version: 1.0

CA397318014 [↗](#)

1031446 (ClinVar) [↗](#)

Gene: GP1BA ([HGNC:2811](#))

Condition: Bernard-Soulier syndrome ([MONDO:0009276](#))

Inheritance Mode: Autosomal recessive inheritance

UUID: 42cd378a-1c69-4e17-921c-a6d05ff77a09

Approved on: 2026/2/5

Published on: 2026/2/5

HGVS expressions

NM_000173.7:c.673T>C

NM_000173.7(GP1BA):c.673T>C (p.Cys225Arg)

NC_000017.11:g.4933277T>C

CM000679.2:g.4933277T>C

NC_000017.10:g.4836572T>C

CM000679.1:g.4836572T>C

NC_000017.9:g.4777352T>C

NG_008767.2:g.5983T>C

ENST00000329125.6:c.673T>C

ENST00000649830.1:c.-888+1065A>G

ENST00000329125.5:c.673T>C

ENST00000611961.1:c.673T>C

NM_000173.6:c.673T>C

Likely Pathogenic

Met criteria codes **6**

PM3_Supporting

PP3

PP4

PM1

PM5

PM2_Supporting

Evidence Links **0**

Expert Panel

[Platelet Disorders VCEP](#) [↗](#)

Criteria Specification Information

[↗](#) **Criteria Specification:** *ClinGen Platelet Disorders Expert*

Panel Specifications to the ACMG/AMP Variant

Interpretation Guidelines for GP1BA Version 1.1.0

[↗](#) **Criteria Specification Approval History**

[↗](#) **Criteria Specifications for this VCEP**

Evidence submitted by expert panel

Platelet Disorders VCEP

The c.673T>C variant in GP1BA is a missense variant predicted to cause substitution of Cysteine by Arginine at amino acid 225 (p.Cys225Arg). At least one patient (internal case from Versiti) with this variant had less than 10% expression of GPIba measured by flow cytometry, which is highly specific for Bernard-Soulier syndrome, the patient also had excessive mucocutaneous bleeding consistent with Bernard-Soulier syndrome (PP4). This individual was compound heterozygous for this variant and the NM_000173.7(GP1BA):c.1601_1602del (p.Tyr534CysfsTer?) variant which was classified as Pathogenic by the PD VCEP. The phase of the

variants in this patient is unknown (0.5 points, PM3_Supporting). This variant resides at a disulfide bond residue, Cys225 GP1BA that is defined as a critical functional domain by the ClinGen PD VCEP (PM1). The largest population MAF allele frequency in gnomAD v4.1.0 is 0.000001695 (based on 2/1179886 alleles) in the European (non-Finnish) population, which is lower than the ClinGen PD VCEP threshold (<0.0001114; PM2_Supporting). The computational predictor REVEL gives a score of 0.653, which is above the ClinGen PD VCEP threshold of >0.644 and predicts a damaging effect on function (PP3). Another missense variant (c.673T>A, p.Cys225Ser) in the same codon has been classified as Pathogenic for Bernard-Soulier syndrome by the ClinGen PD VCEP (PM5). In summary, this variant meets the criteria to be classified as Likely Pathogenic for autosomal recessive Bernard-Soulier syndrome based on the ACMG/AMP criteria applied, as specified by the ClinGen PD VCEP: PP4, PM1, PM5, PM2_Supporting, PM3_supporting and PP3 (VCEP specifications version 1).

Met criteria codes

PM3_Supporting			This variant has been detected in at least 1 proband with Bernard-Soulier syndrome (internal case from Versiti). This individual was compound heterozygous for this variant and the NM_000173.7(GP1BA):c.1601_1602del (p.Tyr534CysfsTer?) variant which was classified as Pathogenic by the PD VCEP. These phase in this variants is unknown (0.5 points, PM3_Supporting).
PP3			The computational predictor REVEL gives a score of 0.653, which is above the ClinGen PD VCEP threshold of >0.644 and predicts a damaging effect on function (PP3).
PP4			At least one patient (internal case from Versiti) with this variant had less than 10% expression of GPIba measured by flow cytometry, which is highly specific for Bernard-Soulier syndrome Additionally, the patient had excessive mucocutaneous bleeding which are consistent with Bernard-Soulier syndrome (PP4).
PM1			This variant resides at a disulfide bond residue, Cys225 GP1BA that is defined as a critical functional domain by the ClinGen PD VCEP (PM1).
PM5			Another missense variant (c.673T>A, p.Cys225Ser) in the same codon has been classified as Pathogenic for Bernard-Soulier syndrome by the ClinGen PD VCEP (PM5).
PM2_Supporting			The largest population MAF allele frequency in gnomAD v4.1.0 is 0.000001695 (based on 2/1179886 alleles) in the European (non-Finnish) population, which is lower than the ClinGen PD VCEP threshold (<0.0001114; PM2_Supporting).

Curation History

▼

▼

Showing 1 to 1 of 1 rows

--

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.