

Variant: *NM_000162.5(GCK):c.56T>A (p.Ile19Asn)*

Version: 1.0

CA367403826 [↗](#)

2136531 (ClinVar) [↗](#)

Gene: GCK ([HGNC:2645](#))

Condition: monogenic diabetes ([MONDO:0015967](#))

Inheritance Mode: Semidominant inheritance

UUID: 588c71a8-9a58-4904-8a95-4da3714631e8

Approved on: 2026/2/24

Published on: 2026/2/26

HGVS expressions

NM_000162.5:c.56T>A

NM_000162.5(GCK):c.56T>A (p.Ile19Asn)

NC_000007.14:g.44153453A>T

CM000669.2:g.44153453A>T

NC_000007.13:g.44193052A>T

CM000669.1:g.44193052A>T

NC_000007.12:g.44159577A>T

NG_008847.1:g.40971T>A

NG_008847.2:g.49718T>A

ENST00000395796.8:c.*54T>A

ENST00000616242.5:c.56T>A

ENST00000682635.1:n.542T>A

ENST00000345378.7:c.59T>A

ENST00000403799.8:c.56T>A

ENST00000671824.1:c.56T>A

ENST00000673284.1:c.56T>A

ENST00000345378.6:c.59T>A

ENST00000395796.7:c.53T>A

ENST00000403799.7:c.56T>A

ENST00000437084.1:c.56T>A

ENST00000476008.1:n.491T>A

ENST00000616242.4:c.53T>A

NM_000162.3:c.56T>A

NM_033507.1:c.59T>A

NM_033508.1:c.53T>A

NM_000162.4:c.56T>A

NM_001354800.1:c.56T>A

NM_033507.2:c.59T>A

NM_033508.2:c.53T>A

NM_033507.3:c.59T>A

NM_033508.3:c.53T>A

Pathogenic

Met criteria codes **7**

PM3

PS4

PM2_Supporting

PP1

PP2

PP3

PP4_Moderate

Expert Panel

Monogenic Diabetes VCEP [↗](#)

Not Met criteria codes 2

PM1

PS3

Evidence Links 0

Criteria Specification Information

[Criteria Specification:](#) *ClinGen Monogenic Diabetes Expert Panel Specifications to the ACMG/AMP Variant Interpretation Guidelines for GCK Version 3.1.0*

[Criteria Specification Approval History](#)

[Criteria Specifications for this VCEP](#)

Evidence submitted by expert panel

Monogenic Diabetes VCEP

The c.56T>A variant in the glucokinase gene, GCK, causes an amino acid change of isoleucine to asparagine at codon 19 (p.(Ile19Asn)) of NM_000162.5. GCK is defined by the ClinGen MDEP as a gene that has a low rate of benign missense variation and has pathogenic missense variants as a common mechanism of disease (PP2). This variant has an incomputable gnomAD v4.1.0 Grpmax filtering allele frequency due to no more than one copy in any subpopulation, thereby meeting the ClinGen MDEP threshold criteria for PM2_Supporting (Grpmax FAF <= 0.000003) (PM2_Supporting). This variant is predicted to be deleterious by computational evidence, with a REVEL score of 0.861, which is greater than the MDEP VCEP threshold of 0.70 (PP3). This variant was identified in 7 unrelated individuals with hyperglycemia (PS4; PMID: 25665835, 33852230, 17573900, internal lab contributors). This variant was identified in an individual with a clinical history highly specific for GCK-hyperglycemia (fasting glucose 5.5-8 mmol/L and HbA1c 5.6 - 7.6% and negative antibody) (PP4_Moderate; internal lab contributors), and segregated with hyperglycemia, with two informative meioses in two families (PP1; PMID: 25665835, internal lab contributors). This variant has been detected in two siblings with neonatal diabetes who were compound heterozygous (confirmed in trans) for this variant and c.1322C>G p.Ser441Trp, which was classified as pathogenic by the MDEP VCEP (PMID: 25665835) (PM3). The relative activity index (RAI) of this variant was above the MDEP cutoff of 0.5 and a relative stability index was not calculated, so PS3 will not be applied (PMID: 25665835). In summary, c.56T>A meets the criteria to be classified as pathogenic for monogenic diabetes. ACMG/AMP criteria applied, as specified by the ClinGen MDEP (specification version 3.1.0, approved 10/10/2025): PS4, PP4_moderate, PP1, PM3, PP2, PP3, PM2_Supporting.

Met criteria codes

PM3			This variant has been detected in two siblings with neonatal diabetes who were compound heterozygous (confirmed in trans) for this variant and c.1322C>G p.Ser441Trp, which was classified as pathogenic by the MDEP VCEP (PMID: 25665835) (PM3).
PS4			This variant was identified in 7 unrelated individuals with hyperglycemia (PS4; PMID: 25665835, 33852230, 17573900, internal lab contributors).
PM2_Supporting			This variant has an incomputable gnomAD v4.1.0 Grpmax filtering allele frequency due to no more than one copy in any subpopulation, thereby meeting the ClinGen MDEP threshold criteria for PM2_Supporting (Grpmax FAF <= 0.000003) (PM2_Supporting).
PP1			This variant segregated with hyperglycemia, with two informative meioses in two families (PP1; PMID: 25665835, internal lab contributors).
PP2			GCK is defined by the ClinGen MDEP as a gene that has a low rate of benign missense variation and has pathogenic missense variants as a common mechanism of disease (PP2).
PP3			

This variant is predicted to be deleterious by computational evidence, with a REVEL score of 0.861, which is greater than the MDEP VCEP threshold of 0.70 (PP3).

PP4_Moderate			This variant was identified in an individual with a clinical history highly specific for GCK-hyperglycemia (fasting glucose 5.5-8 mmol/L and HbA1c 5.6 - 7.6% and negative antibody) (PP4_Moderate; internal lab contributors).
Not Met criteria codes			
PM1			No code specific comments provided, please refer to the summary above or general recommendations provided in the guideline
PS3			The relative activity index (RAI) of this variant was above the MDEP cutoff of 0.5 and a relative stability index was not calculated (PMID: 25665835).

Curation History



Showing 1 to 1 of 1 rows

--

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.