

Variant: NM_000277.2(PAH):c.1A>G (p.Met1Val)

CA114360 [↗](#)

586 (ClinVar) [↗](#)

Gene: PAH ([HGNC:5053](#))

Condition: phenylketonuria ([MONDO:0009861](#))

Inheritance Mode: Autosomal recessive inheritance

UUID: 89f04437-ed5d-4735-8c4a-a9b1d91d10ea

Approved on: 2018-08-07

Published on: 2018-12-10

HGVS expressions

NM_000277.2(PAH):c.1A>G (p.Met1Val)

NM_000277.2:c.1A>G

NC_000012.12:g.102917130T>C

CM000674.2:g.102917130T>C

NC_000012.11:g.103310908T>C

CM000674.1:g.103310908T>C

NC_000012.10:g.101835038T>C

NG_008690.1:g.5473A>G

NG_008690.2:g.46281A>G

NM_000277.1:c.1A>G

NM_001354304.1:c.1A>G

NM_000277.3:c.1A>G

ENST00000307000.7:c.-147A>G

ENST00000546844.1:c.1A>G

ENST00000547319.1:n.312A>G

ENST00000549111.5:n.97A>G

ENST00000551337.5:c.1A>G

ENST00000551988.5:n.90A>G

ENST00000553106.5:c.1A>G

ENST00000635500.1:n.29-4232A>G

Pathogenic

Met criteria codes **4**

PP4_Moderate

PS3

PM2

PM3

Not Met criteria codes **1**

PVS1

Evidence Links **2**

Expert Panel

Phenylketonuria VCEP [↗](#)

Criteria Specification Information **!**

[↗](#) **Criteria Specifications for this VCEP**

Evidence submitted by expert panel

Phenylketonuria VCEP

PAH-specific ACMG/AMP criteria applied: PM2: gnomAD MAF=0.00002; PP4_Moderate: Seen in PKU patients. BH4 disorders ruled out. (PMID:2574002); PS3: <3% (PMID:9450897). PM3: Detected in trans with known pathogenic variants. In summary this variant meets criteria to be classified as pathogenic for phenylketonuria in an autosomal recessive manner based on the ACMG/AMP criteria applied as specified by the PAH Expert Panel: (PM2, PM3, PP4_Moderate, PS3). Updated to reflect new PVS1 recommendations.

Met criteria codes

PP4_Moderate			Seen in PKU patients. BH4 disorders ruled out.
Analysis of RFLP haplotypes and mutations revealed a novel mutation, an A-to-G transition (met----val) in codon 1 (the translation-initiation codon). It occurred on 5 of the 18 mutant chromosomes. A proband homozygous for this mutation had the PKU phenotype. The appropriate biochemical tests were done to rule out disorders of tetrahydrobiopterin homeostasis. PubMed:2574002 			
PS3			<3%
< 3 PAH enzyme activity as % of wild type. (p91023(B)/COS PubMed:9450897 			
PM2			gnomAD MAF=0.00002
PM3			Detected in trans with c.1315+1G>A (known pathogenic), and also in the homozygous state.
A proband was homozygous for this mutation (A-to-G transition (met-val) in codon1]. In other probands, the codon1 mutation was inherited once with the splice junction mutation in exon 12 (c.1315+1G>A),and was inherited twice with a mutation on haplotype 1. PubMed:2574002 			

Not Met criteria codes

PVS1			Initiation codon variant
------	---	---	--------------------------

Curation History



 

 

Showing 1 to 1 of 1 rows

--

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.

[ClinGen Terms of Use.](#)

[✕ Powered by BCM's Genboree.](#)