

Variant: *NM_000260.4(MYO7A):c.6062A>G (p.Lys2021Arg)*

Version: 2.0

CA10576905 [↗](#)

228282 (ClinVar) [↗](#)

Gene: [MYO7A](#)

Condition: Usher syndrome type 1 ([MONDO:0010168](#))

Inheritance Mode: Autosomal recessive inheritance

UUID: c7b98a09-bc39-416b-af89-f1dfff058317

Approved on: 2023/1/18

Published on: 2023/2/6

HGVS expressions

NM_000260.4:c.6062A>G

NM_000260.4(MYO7A):c.6062A>G (p.Lys2021Arg)

NM_000260.4(MYO7A):c.6062A>G

NC_000011.10:g.77211162A>G

CM000673.2:g.77211162A>G

NC_000011.9:g.76922207A>G

CM000673.1:g.76922207A>G

NC_000011.8:g.76599855A>G

NG_009086.1:g.87898A>G

NG_009086.2:g.87917A>G

ENST00000409709.9:c.6062A>G

ENST00000670577.1:c.3863A>G

ENST00000409619.6:c.5915A>G

ENST00000409709.7:c.6062A>G

ENST00000458169.2:c.3488A>G

ENST00000458637.6:c.5948A>G

ENST00000481328.7:n.3598A>G

ENST00000526863.2:n.25+251A>G

ENST00000605744.1:n.1529A>G

NM_000260.3:c.6062A>G

NM_001127180.1:c.5948A>G

NM_001127180.2:c.5948A>G

NM_001369365.1:c.5915A>G

Likely Pathogenic

Met criteria codes **4**

PP3 PP4 PM3 PM2_Supporting

Evidence Links **0**

Expert Panel

[Hearing Loss VCEP](#) [↗](#)

Criteria Specification Information

[↗](#) **Criteria Specification:** *ClinGen Hearing Loss Expert Panel Specifications to the ACMG/AMP Variant Interpretation Guidelines for CDH23, COCH, GJB2, KCNQ4, MYO6, MYO7A, SLC26A4, TECTA and USH2A Version 2*

[↗](#) **Criteria Specification Approval History**

[↗](#) **Criteria Specifications for this VCEP**

Hearing Loss VCEP

The c.6062A>G (NM_000260.4(MYO7A):c.6062A>G (p.Lys2021Arg)) variant in MYO7A is a missense variant predicted to cause substitution of lysine by arginine at amino acid 2021. This variant is absent from gnomAD v2.1.1 (PM2_Supporting). The computational predictor REVEL gives a score of 0.748, which is above the threshold of 0.7, evidence that correlates with impact to MYO7A function (PP3). This variant has been detected in at least 4 individuals with Usher Syndrome type 1 (from three families). For one of those individuals, they were compound heterozygous for the variant and a pathogenic/likely pathogenic variant (phase unknown) ((NM_000260.4(MYO7A):c.722G>A (p.Arg241His); 0.5 PM3 points; 21436283). In another family, this variant was found in the homozygous state in a male adolescent, 14 years of age, with bilateral congenital hearing loss and features of night blindness (0.5 PM3 points; Allan, 2014; 1). Due to consanguinity, 2 individuals from the same family were homozygous for the variant (0.25 PM3 points; Allam, 2014; 1) (PM3). At least one patient with this disease displayed sensorineural hearing loss and retinitis pigmentosa, which is highly specific for Usher Syndrome 1 (PMID: 21436283; PP4). This variant was re-reviewed on 1.18.2023 and because no additional evidence is available, professional judgment was used to retain this variant as likely pathogenic for autosomal recessive Usher Syndrome based on the ACMG/AMP criteria applied, as specified by the ClinGen Hearing Loss VCEP (PM2_supporting, PP3, PM3, PP4; Version 2; 2022). 1. https://docs.google.com/document/d/17VJ_cidV8dqRQMHTzSVrFa-7PrmJ4trkw2wQsNEPqv/edit

Met criteria codes

PP3	 	The REVEL score 0.748 (above the threshold for pathogenicity set by the Hearing Loss VCEP). The variant is very well-conserved in UCSC database across species, except for 1 mammal (elephant) which has Thr at this site. The results from 3 in silico predictors/lines of evidence, [PolyPhen2, conservation, REVEL score], provide evidence that correlates with impact to MYO7A function (PP3).
PP4	 	All patients evaluated had hearing loss, which is highly specific for Usher Syndrome Type 1 (PP4).
PM3	 	Variant was reported in an individual from France with Usher Syndrome 1. This variant was found in the compound heterozygous state with another variant (NM_000260.4(MYO7A):c.722G>A p.Arg241His), which is pathogenic/likely pathogenic. Phase was not reported (0.5 points; PM3; PMID: 21436283). Variant was found in the homozygous state at least two children (boy age nine and girl age thirteen) from the same family, each of whom had sensorineural hearing loss with early symptoms of retinal degeneration. The parents were consanguineous (0.25 points; PM3; Allam, 2014). In another family, this variant was found in the homozygous state in a male adolescent, 14 years of age, with bilateral congenital hearing loss, trouble seeing at night, and chronic pain in the throat (0.5 points; PM3; Allam, 2014).
PM2_Supporting	 	This variant is absent from gnomAD v2.1.1 (PM2_Supporting).

▾

▾

Showing 1 to 2 of 2 rows

--

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.