

Variant: NM_000527.5(LDLR):c.1003G>T (p.Gly335Cys)

Version: 1.0

CA404082719 [↗](#)

1395739 (ClinVar) [↗](#)

Gene: LDLR ([HGNC:3949](#))

Condition: hypercholesterolemia, familial ([MONDO:0007750](#))

Inheritance Mode: Semidominant inheritance

UUID: f40ccd74-b0bc-4f96-bbe1-ad5342f0c802

Approved on: 2023/4/28

Published on: 2023/5/1

HGVS expressions

NM_000527.5:c.1003G>T

NM_000527.5(LDLR):c.1003G>T (p.Gly335Cys)

NC_000019.10:g.11110714G>T

CM000681.2:g.11110714G>T

NC_000019.9:g.11221390G>T

CM000681.1:g.11221390G>T

NC_000019.8:g.11082390G>T

NG_009060.1:g.26334G>T

ENST00000252444.10:c.1261G>T

ENST00000559340.2:c.1003G>T

ENST00000560467.2:c.941-800G>T

ENST00000558518.6:c.1003G>T

ENST00000252444.9:c.1257G>T

ENST00000455727.6:c.499G>T

ENST00000535915.5:c.880G>T

ENST00000545707.5:c.622G>T

ENST00000557933.5:c.1003G>T

ENST00000558013.5:c.1003G>T

ENST00000558518.5:c.1003G>T

ENST00000560173.1:n.2G>T

ENST00000560467.1:c.541-800G>T

NM_000527.4:c.1003G>T

NM_001195798.1:c.1003G>T

NM_001195799.1:c.880G>T

NM_001195800.1:c.499G>T

NM_001195803.1:c.622G>T

NM_001195798.2:c.1003G>T

NM_001195799.2:c.880G>T

NM_001195800.2:c.499G>T

NM_001195803.2:c.622G>T

Likely Pathogenic

Met criteria codes 4

PS3_Moderate PP3 PP4 PM2

Not Met criteria codes 22

BP5 BP7 BP4 BP3 BP1 BP2
PS1 PS2 PS4 PP1 PP2 PM6
PM1 PM3 PM5 PM4 PVS1
BA1 BS2 BS1 BS4 BS3

Evidence Links 1

Expert Panel

Familial Hypercholesterolemia VCEP

Criteria Specification Information

- Criteria Specification: ClinGen Familial Hypercholesterolemia Expert Panel Specifications to the ACMG/AMP Variant Classification Guidelines Version 1.2
- PDF
- Criteria Specification Approval History
- Criteria Specifications for this VCEP

Evidence submitted by expert panel

Familial Hypercholesterolemia VCEP

The NM_000527.5(LDLR):c.1003G>T (p.Gly335Cys) variant is classified as Likely pathogenic for Familial Hypercholesterolemia by applying evidence codes PM2, PS3_Moderate, PP3 and PP4 as defined by the ClinGen Familial Hypercholesterolemia Expert Panel LDLR-specific variant curation guidelines (https://doi.org/10.1016/j.gim.2021.09.012). The supporting evidence is as follows: PM2 - This variant is absent from gnomAD (gnomAD v2.1.1). PS3_moderate - Level 2 FS: Hu et al. 2021 (PMID: 34970301): Heterologous cells (HEK293), Western blot and Dil-LDL (just uptake) assays - results: less mature LDLR detected, 43% uptake. ---- part of the LDLR cycle (uptake) is less than 70% activity of wild-type, so PS3_Moderate is met. PP3 - REVEL = 0.889. It is above 0.75, so PP3 is met. PP4 - variant meets PM2 and was identified in 1 index case from Hu et al. 2021 (PMID: 34970301) with 5.5mmol/L LDL under no medication, and family history of hypercholesterolemia, so Simon-Broome possible is met; so PP4 is met.

Met criteria codes

PS3_Moderate			Level 2 FS: Hu et al. 2021 (PMID: 34970301): Heterologous cells (HEK293), Western blot and Dil-LDL (just uptake) assays - results: less mature LDLR detected, 43% uptake. ---- part of the LDLR cycle (uptake) is less than 70% activity of wild-type, so PS3_Moderate is met
			Heterologous cells (HEK293), Western blot and Dil-LDL (just uptake) assays - results: less mature LDLR detected, 43% uptake. PubMed:34970301
PP3			REVEL = 0.889. It is above 0.75, so PP3 is met
PP4			variant meets PM2 and was identified in - 2 unrelated index cases from Leren et al. 2021 (PMID: 33740630), but inclusion criteria was "No stringent criteria have been used to select hypercholesterolemic subjects for genetic testing, except that a value for total serum cholesterol of at least 6 mmol/l has been required for molecular genetic testing in adults.", which is not enough to insure clinical FH criteria, so not considered; - 1 index case from Hu et al. 2021 (PMID: 34970301) with 5.5mmol/L LDL under no medication, and family history of hypercholesterolemia, so Simon-Broome possible is met; so PP4 is met
PM2			This variant is absent from gnomAD (gnomAD v2.1.1).

Not Met criteria codes

BP5			not applicable
BP7			variant is missense, so not met
BP4			REVEL = 0.889. It is not below 0.50, so BP4 is not met
BP3			not applicable
BP1			not applicable
BP2			variant identified in 1 index case from Hu et al. 2021 (PMID: 34970301) who is also heterozygous (confirmed in trans) for LDLR c.226G>C, p.Gly76Arg and has LDL of 5.62mmol/L under atorvastatin 20mg. Calculated untreated LDL: 5.62 / 0.57 (Ruel et al., 2018) = 9.86mmol/L, which is not clearly a homozygous phenotype (>13mmol/L) nor a clearly heterozygous phenotype (<8mmol/L), so not met.
PS1			There are no other variants that lead to the same amino acid change, so not met
PS2			there is no data on de novo occurrence
PS4			variant meets PM2 and was identified in - 2 unrelated index cases from Leren et al. 2021 (PMID: 33740630), but inclusion criteria was "No stringent criteria have been used to select hypercholesterolemic subjects for genetic testing, except that a value for total serum cholesterol of at least 6 mmol/l has been required for molecular genetic testing in adults.", which is not enough to insure clinical FH criteria, so not considered; - 1 index case from Hu et al. 2021 (PMID: 34970301) with 5.5mmol/L LDL under no medication, and family history of hypercholesterolemia, so Simon-Broome possible is met; ---- 1 index case, not enough, so not met
PP1			variant identified in - 2 relatives from Leren et al. 2021 (PMID: 33740630), but there is no data on phenotype, so not considered; - 1 family from Hu et al. 2021 (PMID: 34970301), but there are 2 variants in the family, so not considered.
PP2			not applicable
PM6			there is no data on de novo occurrence
PM1			variant is missense and meets PM2, but it is not in exon 4 and does not alter Cys, so not met
PM3			variant identified in 1 index case from Hu et al. 2021 (PMID: 34970301) who is also heterozygous (confirmed in trans) for LDLR c.226G>C, p.Gly76Arg and has LDL of 5.62mmol/L under atorvastatin 20mg. Calculated untreated LDL: 5.62 / 0.57 (Ruel et al., 2018) = 9.86mmol/L, which is not clearly a homozygous phenotype (>13mmol/L) nor a clearly heterozygous phenotype (<8mmol/L), so not met.
PM5			There are 4 other missense variants in the same codon: - NM_000527.5(LDLR):c.1003G>A (p.Gly335Ser) ClinVar ID 183105 - classified as Likely pathogenic by these guidelines - NM_000527.5(LDLR):c.1003G>C (p.Gly335Arg) ClinVar ID 926520 - classified as VUS by these guidelines - NM_000527.5(LDLR):c.1004G>A (p.Gly335Asp) ClinVar ID 225181

- classified as VUS by these guidelines - NM_000527.5(LDLR):c.1004G>T (p.Gly335Val) ClinVar ID 251587 - classified as VUS by these guidelines no variant classified as Pathogenic, so not met

PM4			variant is missense, so not met
PVS1			variant is missense and not in initiation codon, so not met
BA1			This variant is absent from gnomAD (gnomAD v2.1.1).
BS2			Variant has not been identified in normolipidemic individuals, so not met
BS1			This variant is absent from gnomAD (gnomAD v2.1.1).
BS4			variant identified in - 2 relatives from Leren et al. 2021 (PMID: 33740630), but there is no data on phenotype, so not considered; - 1 family from Hu et al. 2021 (PMID: 34970301), but there are 2 variants in the family, so not considered.
BS3			Level 2 FS: Hu et al. 2021 (PMID: 34970301): Heterologous cells (HEK293), Western blot and Dil-LDL (just uptake) assays - results: less mature LDLR detected, 43% uptake. ---- part of the LDLR cycle (uptake) is less than 70% activity of wild-type, so BS3 is not met

Curation History

▼

▼

Showing 1 to 1 of 1 rows

--

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.