

Variant: *NM_000330.4(RS1):c.655T>C (p.Cys219Arg)*

Version: 1.0

CA226841 [↗](#)

99015 (ClinVar) [↗](#)

Gene: RS1 ([HGNC:6247](#))

Condition: X-linked retinoschisis ([MONDO:0010725](#))

Inheritance Mode: X-linked inheritance (recessive (HP:0001419))

UUID: f559e124-d1f2-4d0e-b199-d15282691f6e

Approved on: 2025/10/4

Published on: 2025/10/5

HGVS expressions

NM_000330.4(RS1):c.655T>C (p.Cys219Arg)

NM_000330.4:c.655T>C

NC_000023.11:g.18642024A>G

CM000685.2:g.18642024A>G

NC_000023.10:g.18660144A>G

CM000685.1:g.18660144A>G

NC_000023.9:g.18570065A>G

NG_008475.1:g.221420A>G

NG_008659.3:g.40425T>C

ENST00000379984.4:c.655T>C

ENST00000379984.3:c.655T>C

ENST00000379989.6:c.2714-3983A>G

ENST00000379996.7:c.2714-3983A>G

ENST00000476595.1:n.1146T>C

NM_000330.3:c.655T>C

NM_001037343.1:c.2714-3983A>G

NM_003159.2:c.2714-3983A>G

NM_001037343.2:c.2714-3983A>G

NM_003159.3:c.2714-3983A>G

Likely Pathogenic

Met criteria codes **6**

PS3_Supporting

PP3_Strong

PP1

PM5_Supporting

PS4_Supporting

PM2_Supporting

Not Met criteria codes **3**

PM6

PM1

PS2

Evidence Links **0**

Expert Panel

X-linked Inherited Retinal Disease VCEP [↗](#)

Criteria Specification Information

[↗](#) **Criteria Specification:** ClinGen X-linked Inherited Retinal Disease Expert Panel Specifications to the ACMG/AMP Variant Interpretation Guidelines for RS1 Version 1.0.0

[↗](#) **Criteria Specification Approval History**

[↗](#) **Criteria Specifications for this VCEP**

Evidence submitted by expert panel

X-linked Inherited Retinal Disease VCEP

NM_000330.4(RS1):c.655T>C (p.Cys219Arg) is a missense variant encoding the substitution of cysteine with arginine at amino acid 219. Another missense variant in the same codon, NM_000330.4(RS1):c.655T>G (p.Cys219Gly) has been classified as likely pathogenic for X-linked retinoschisis by the ClinGen X-linked IRD VCEP, while no benign missense variants have been identified in this codon. The present variant has a higher Grantham’s distance (180) than the comparison variant (159), and SpliceAI has been used to confirm that neither variant has a predicted impact on RS1 splicing (PM5_Supporting). This variant is absent from hemizygous individuals in gnomAD v4.1.0 (PM2_Supporting). This variant has been reported in at least 2 unrelated probands meeting the PS4 requirement of a male diagnosed with X-linked retinoschisis (PMID: 39462066, PMID: 35982512, PMID: 9618178, PS4_Supporting). The variant has been reported to segregate with X-linked retinoschisis through 2 affected family members (PP1; PMID: 39462066). This variant has been identified as a de novo occurrence with unconfirmed parental relationships in 1 individual meeting the PS2 phenotype requirement of a male diagnosed with retinoschisis (PMID: 9618178). No description was provided of parental genotyping, so PS2 was not met. COS-1 cells exogenously expressing the variant protein exhibit loss of RS1 secretion into the medium relative to the wild-type control, caused by misfolding, aggregation, and ER retention of the variant protein (PMID: 12746437, PS3_Supporting). The computational predictor REVEL gives a score of 0.978, which is above the ClinGen X-linked IRD VCEP threshold of >0.932 and predicts a damaging effect on RS1 function (PP3_Strong). The computational splicing predictor SpliceAI gives a delta score of 0.00, which is below the ClinGen X-linked IRD VCEP threshold of ≥0.2 and does not predict that the variant disrupts RS1 splicing. This variant is a missense substitution affecting a critical amino acid residue involved in disulfide bridge formation as defined by the ClinGen X-linked IRD VCEP (PMID: 26812435). PM1_Strong is not met as this code is ineligible to be used in combination with PP3_Strong. In summary, this variant is classified as likely pathogenic for X-linked retinoschisis based on the ClinGen X-linked Inherited Retinal Disease Expert Panel Specifications to the ACMG/AMP Variant Interpretation Guidelines for RS1 Version 1.0.0: PM2_Supporting, PP3_Strong, PM5_Supporting, PS3_Supporting, PS4_Supporting, and PP1.

Met criteria codes

PS3_Supporting			COS-1 cells exogenously expressing the variant protein exhibit loss of RS1 secretion into the medium relative to the wild-type control, caused by misfolding, aggregation, and ER retention of the variant protein (PMID: 12746437, PS3_Supporting).
PP3_Strong			The computational predictor REVEL gives a score of 0.978, which is above the ClinGen X-linked IRD VCEP threshold of >0.932 and predicts a damaging effect on RS1 function (PP3_Strong). The computational splicing predictor SpliceAI gives a delta score of 0.00, which is below the ClinGen X-linked IRD VCEP threshold of ≥0.2 and does not predict that the variant disrupts RS1 splicing.
PP1			The variant has been reported to segregate with X-linked retinoschisis through 2 affected family members (PP1; PMID: 39462066)
PM5_Supporting			Another missense variant in the same codon, NM_000330.4(RS1):c.655T>G (p.Cys219Gly) has been classified as likely pathogenic for X-linked retinoschisis by the ClinGen X-linked IRD VCEP, while no benign missense variants have been identified in this codon. The present variant has a higher Grantham’s distance (180) than the comparison variant (159), and SpliceAI has been used to confirm that neither variant has a predicted impact on RS1 splicing (PM5_Supporting).
PS4_Supporting			This variant has been reported in at least 2 unrelated probands meeting the PS4 requirement of a male diagnosed with XLR5 (PMIDs: 39462066, 35982512, 9618178, PS4_supporting).
PM2_Supporting			This variant is absent from hemizygous individuals in gnomAD v4.1.0 (PM2_Supporting).

Not Met criteria codes			
PM6		✖	No code specific comments provided, please refer to the summary above or general recommendations provided in the guideline
PM1		✖	This variant is a missense substitution affecting a critical amino acid residue involved in disulfide bridge formation as defined by the ClinGen X-linked IRD VCEP (PMID: 26812435). PM1_Strong is not met as this code is ineligible to be used in combination with PP3_Strong.
PS2		✖	This variant has been identified as a de novo occurrence with unconfirmed parental relationships in 1 individual meeting the PS2 phenotype requirement of a male diagnosed with retinoschisis (PMID: 9618178). No description was provided of parental genotyping, so PS2 was not met.

Curation History 		
<div><div></div><div>▼</div><div>▼</div></div>		
Showing 1 to 1 of 1 rows		
<div></div>		

The information on this website is not intended for direct diagnostic use or medical decision-making without review by a genetics professional. Individuals should not change their health behavior solely on the basis of information contained on this website. If you have questions about the information contained on this website, please see a health care professional.